

UCMC Transplant Program - Infectious Prophylaxis Guidelines

For the following recipients: (1) Pancreas Alone, (2) Pancreas after Kidney, and (3) Simultaneous Pancreas-Kidney

	Patient Population		Medication	Dosing Regimen	
Peri-operative	All non-penicillin-allergic		Ceftriaxone	Pre-op ceftriaxone: 2 gm IV on call to OR Post-op ceftriaxone: 2 gm IV x 1 dose, given 24 hours after pre-op dose	
	Penicillin-allergic		Vancomycin AND ciprofloxacin	Pre-op:	<ul style="list-style-type: none"> • Vancomycin: 20 mg/kg IV on call to OR • Ciprofloxacin: ≤ 80 kg: 400 mg IV on call to OR; > 80 kg: 600 mg IV on call to OR
				Post-op:	<ul style="list-style-type: none"> • Vancomycin: 15 mg/kg IV x 1 dose, given 12 hours after pre-op dose; then dosed per renal function for 48-hour total duration • Ciprofloxacin: 400 mg IV x 1 dose, given 12 hours after pre-op dose; then dosed per renal function for 48-hour total duration
All		Fluconazole ¹	200 mg IV x 1 dose	1 dose	
Fungal Initiate POD1	All		Fluconazole ¹	200 mg PO daily	1 month
PJP Initiate POD 1-2	All non-sulfa-allergic		Bactrim SS ²	1 tablet PO Monday, Wednesday, Friday	6 months (lifelong in HIV+ recipients)
	Sulfa-allergic		Dapsone ³ Pentamidine ⁴ Atovaquone	100 mg PO daily 300 mg by nebulization once monthly 1500 mg PO daily	6 months (lifelong in HIV+ recipients)
CMV Initiate POD 1-2 <i>If concern for renal function or marrow suppression adjust dose as described below</i>	High risk	CMV IgG Donor + / Recipient -	Valganciclovir ^{5,6}	900 mg PO daily ⁷	6 months (monitoring ⁸)
	Intermediate risk	CMV IgG Donor + / Recipient + CMV IgG Donor - / Recipient +	Valganciclovir ^{5,6}	900 mg PO daily ⁷	3 months (monitoring ⁸)
	Low risk	CMV IgG Donor - / Recipient -	Acyclovir ⁵	800 mg PO twice daily ⁷	Depends on EBV serostatus ⁹

¹ **Fluconazole dose adjustments:** CrCl < 50 = 100 mg PO daily; CVVH or CVVHD = no dose adjustment necessary
HD = 200mg PO 3x week; administer after each dialysis session on dialysis days only

² **Bactrim (trimethoprim-sulfamethoxazole) SS (single strength) dose adjustments:**

- HD = 1 tablet PO 3x weekly after HD (dialysis days only)
- CVVH or CVVHD = no dose adjustment necessary
- Leukopenia = refer to leukopenia management guidelines

³ **Dapsone:** do not check G6PD routinely; only in those of Mediterranean descent

⁴ **Pentamidine:** premedicate with albuterol 2.5 mg by nebulization.

⁵ **Anti-viral dose adjustments:** ONLY for renal dysfunction (refer to table for dose adjustments)

- LEUKOPENIA = refer to leukopenia management guidelines.
- IF PERSISTENT NEUTROPENIA: hold & monitor CMV RT Quant PCR weekly.
- LETERMОВIR: 2nd option for persistent neutropenia. Consider letermovir pending insurance and nonformulary approval and consult Txp ID. Letermovir use may require initiation of acyclovir for HSV prophylaxis if < POD #30 or undergoing rejection treatment and CMV PCR monitoring every 2 weeks.

⁶ If unable to take PO valganciclovir: convert to ganciclovir 5 mg/kg IV day (adjust for renal function)

⁹ Acyclovir Duration

EBV	EBV IgG	Acyclovir Duration
High risk	Donor + / Recipient -	3 months
Intermediate risk or Low risk	Donor + / Recipient + Donor - / Recipient + Donor - / Recipient -	1 month

⁷ Anti-viral renal dose adjustments:

CrCl (mL/min)	Valganciclovir PO	Ganciclovir IV	Acyclovir PO
>70	900 mg daily	5 mg/kg q 24 hour	800 mg 2x day
60-69	900 mg daily	2.5 mg/kg q 24 hours	800 mg 2x day
50-59	450 mg daily	2.5 mg/kg q 24 hours	800 mg 2x day
40-49	450 mg daily	1.25 mg/kg q 24 hours	800 mg 2x day
25-39	450 mg M-W-F	1.25 mg/kg q 24 hours	800 mg 2x day
10-24	450 mg twice weekly	0.625 mg/kg q 24 hours	400 mg 2x day
<10 or iHD	450 mg twice weekly after iHD	0.625 mg/kg 3x/week after iHD	400 mg 2x day after HD
PD	450 mg twice weekly	0.625 mg/kg 3x/week	400 mg 2x day
CVVH	450 mg q 48 hours	1.25 mg/kg q 24 hours	800 mg 2x day
CVVHD/HDF	450 mg daily	2.5 mg/kg q 24 hours	800 mg 2x day

⁸ CMV monitoring (valganciclovir):

- If prophylaxis is held/delayed: monitor CMV RT Quant PCR weekly until resumed or initiated
- After completion of prophylaxis therapy: monitor CMV RT Quant PCR every 2 weeks x 3. If CMV viremia develops change CMV PCR monitoring to weekly (refer to CMV treatment guidelines)

